

# ON A CONJECTURE OF SHOKUROV: CHARACTERIZATION OF TORIC VARIETIES

YURI G. PROKHOROV

**ABSTRACT.** We verify a special case of V. V. Shokurov's conjecture about characterization of toric varieties. More precisely, let  $(X, D = \sum d_i D_i)$  be a three-dimensional log variety such that  $K_X + D$  is numerically trivial and  $(X, D)$  has only purely log terminal singularities. In this situation we prove the inequality

$$\sum d_i \leq \text{rk Weil}(X)/(\text{algebraic equivalence}) + \dim(X).$$

We describe such pairs for which the equality holds and show that all of them are toric.

## 1. INTRODUCTION

The aim of this note is to discuss the birational characterization of toric varieties. Let  $X$  be a normal projective toric variety and let  $D = \sum_{i=1}^r D_i$  be the sum of invariant divisors. It is well known that the pair  $(X, D)$  has only log canonical singularities (see e. g. [3, 3.7]),  $K_X + D$  is linearly trivial and  $r = \text{rk}(\text{Weil}(X)/\approx) + \dim(X)$ , where  $\text{Weil}(X)$  is the group of Weil divisors and  $\approx$  is the algebraic equivalence.

Shokurov observed that this property can characterize toric varieties:

**Conjecture 1.1** ([12]). *Let  $(X, D = \sum d_i D_i)$  be a projective log variety such that  $(X, D)$  has only log canonical singularities and numerically trivial. Then  $\sum d_i \leq \text{rk}(\text{Weil}(X)/\approx) + \dim(X)$ . Moreover, if the equality holds, then  $(X, \lfloor D \rfloor)$  is a toric pair.*

Shokurov also conjectured the relative version of 1.1 (cf. Theorem 2.3) and he expects that one can replace numerical triviality of  $K_X + D$  with nefness of  $-(K_X + D)$ . We do not discuss these details here.

Conjecture 1.1 was proved in dimension two in [12] (see also [9, Sect. 8] and Proposition 2.1 below). Our main result is the following partial answer on Conjecture 1.1 in dimension three:

**Theorem 1.2.** *Let  $(X, D = \sum d_i D_i)$  be a three-dimensional projective variety over  $\mathbb{C}$  such that  $K_X + D \equiv 0$  and  $(X, D)$  has only purely log terminal singularities. Then*

$$(1.3) \quad \sum d_i \leq \text{rk}(\text{Weil}(X)/\approx) + 3.$$

Moreover, if the equality holds, then up to isomorphisms one of the following holds:

- (i)  $X \simeq \mathbb{P}^3$ ,  $[D] = 0$  or  $[D] = \mathbb{P}^2$ ;
- (ii)  $X \simeq \mathbb{P}^1 \times \mathbb{P}^2$ ,  $[D] = 0$  or  $[D] = \{\text{pt}\} \times \mathbb{P}^2$  or  $[D] = \mathbb{P}^1 \times \{\text{line}\}$ ;
- (iii)  $X \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(d))$ ,  $d \geq 1$ ,  $[D]$  is the section corresponding to the surjection  $\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(d) \rightarrow \mathcal{O}_{\mathbb{P}^1}(d)$ ;
- (iv)  $X \simeq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ ,  $[D] = 0$  or  $[D] = \{\text{pt}\} \times \mathbb{P}^1 \times \mathbb{P}^1$  or  $[D] = \{\text{pt}_1, \text{pt}_2\} \times \mathbb{P}^1 \times \mathbb{P}^1$ ;
- (v)  $X \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(d))$ ,  $d \geq 1$ ,  $[D]$  is the negative section, or a disjoint union of two sections, one of them is negative;
- (vi)  $X \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1} \oplus \mathcal{L})$ ,  $\mathcal{L} \in \text{Pic}(\mathbb{P}^1 \times \mathbb{P}^1)$ ,  $[D]$  is the negative section, or a disjoint union of two sections, one of them is negative.

In all cases  $(X, [D])$  is toric.

Clearly, our theorem is not a characterization of toric varieties, but we hope that Conjecture 1.1 can be proved in a similar way.

This paper was subject of my talk given at Waseda seminar on January 18, 2000. I am grateful to the participants of this seminar, especially Professor S. Ishii, for attention and valuable discussions. I also would like to thank the Department of Mathematics of Tokyo Institute of Technology for hospitality during my stay on 1999–2000. This work was partially supported by the grant INTAS-OPEN-97-2072.

## 2. PRELIMINARIES

**Notation.** All varieties are defined over  $\mathbb{C}$ . Basically we employ the standard notation of the Minimal Model Program (MMP, for short). Throughout this paper  $\rho(X)$  is the Picard number and  $\overline{NE}(X)$  is the Mori cone of  $X$ . We call a pair  $(X, D)$  consisting of a normal algebraic variety  $X$  and a boundary  $D$  on  $X$  a *log variety* or a *log pair*. Here a *boundary* is a  $\mathbb{Q}$ -Weil divisor  $D = \sum d_i D_i$  such that  $0 \leq d_i \leq 1$  for all  $i$ . A *contraction* is a projective morphism  $\varphi: X \rightarrow Z$  of normal varieties such that  $\varphi_* \mathcal{O}_X = \mathcal{O}_Z$ . Abbreviations klt, plt, lc are reserved for Kawamata log terminal, purely log terminal and log canonical, respectively (refer to [11], [4] and [3] for the definitions). Let  $(X, D)$  be a log pair and let  $S := [D]$ . For simplicity, assume that  $(X, D)$  is lc in codimension two. The Adjunction Formula proposed by Shokurov [11, Sect. 3] states  $(K_X + D)|_S = K_S + \text{Diff}_S(D - S)$ , where  $\text{Diff}_S(D - S)$  is a naturally defined effective  $\mathbb{Q}$ -Weil divisor on  $S$ , so-called *different*. Moreover,  $K_X + D$  is plt near  $S$  if and only if  $S$  is normal and  $K_S + \text{Diff}_S(D - S)$  is klt [4, 17.6].  $LCS(X, D)$  denotes the *locus of log canonical singularities* of  $(X, D)$  that is the set of all points where  $(X, D)$  is not klt [11]. Let  $\varphi: X \rightarrow Z$  be any fiber type contraction and let  $D = \sum d_i D_i$  be a  $\mathbb{Q}$ -divisor on  $X$ . We will write  $D = \sum_{\text{ver}} d_i D_i + \sum_{\text{hor}} d_i D_i = D^{\text{ver}} + D^{\text{hor}}$ , where  $\sum_{\text{ver}}$  (respectively  $\sum_{\text{hor}}$ )

runs through all components  $D_i$  such that  $\dim \varphi(D_i) < \dim(D_i)$  (respectively  $\varphi(D_i) = Z$ ). We will frequently use the above notation without reference.

In dimension two Conjecture 1.1 is much easier than higher dimensional one. We need only the following weaker version:

**Proposition 2.1.** *Let  $(X, D = \sum d_i D_i)$  be a projective log surface such that  $-(K_X + D)$  is nef and  $(X, D)$  is lc. Then  $\sum d_i \leq \rho(X) + 2$ . Moreover, if the equality holds and  $(X, D)$  is klt, then  $X \simeq \mathbb{P}^2$ , or  $X \simeq \mathbb{P}^1 \times \mathbb{P}^1$ .*

For the general statement we refer to [12], see also [9].

*Proof.* Assume that  $\sum d_i - \rho(X) - 2 \geq 0$  and run  $K_X$ -MMP. According to [1], Log MMP works even in the category of log canonical pairs. On each step,  $\sum d_i - \rho(X) - 2$  does not decrease and all assumptions are preserved (see [4, 2.28]). At the end we get one of the following:

**Case 1.**  $\rho(X) = 1$ . Then  $\sum d_i - \rho(X) - 2 \leq 0$  by [4, 18.24].

**Case 2.** There is an extremal contraction onto a curve  $\varphi: X \rightarrow Z$  (in particular,  $\rho(X) = 2$ ). Let  $\ell$  be a general fiber. Then

$$(2.2) \quad 2 = -K_X \cdot \ell \geq D \cdot \ell \geq \sum_{\text{hor}} d_i.$$

Hence  $\sum_{\text{ver}} d_i \geq 2$  and  $K_X + D^{\text{hor}}$  is not nef. Let  $\phi: X \rightarrow W$  is a contraction of  $(K + D^{\text{hor}})$ -negative extremal ray. If  $\phi$  is birational, we replace  $X$  with  $W$  and obtain Case 1 above. Thus we may assume that  $W$  is a curve, so  $\varphi$  and  $\phi$  are symmetric. As above,  $\sum_{\text{ver}} d_i \leq 2$ , so  $\sum d_i = 4$ .

Now assume that  $(X, D)$  is klt and  $\sum d_i - \rho(X) - 2 = 0$ . Then after each divisorial contraction  $\sum d_i - \rho(X) - 2$  increases. Hence we have only cases 1 or 2 above. In Case 1,  $X \simeq \mathbb{P}^2$  by Lemma 3.1 below. In Case 2 we have the equality in (2.2), so  $D_i \cdot \phi^{-1}(w) = 1$  for any component of  $D^{\text{ver}}$  and a general fiber  $\phi^{-1}(w)$ . Therefore  $D_i$  is not a multiple fiber of  $\varphi$  and  $X$  is smooth along  $D_i$ . Similarly, if  $D_j$  is a component of  $D^{\text{hor}}$ , then  $\phi(D_j) = \text{pt}$ ,  $X$  is smooth along  $D_j$  and  $D_i \cdot D_j = 1$ . Finally,  $\varphi \times \phi: X \rightarrow Z \times W = \mathbb{P}^1 \times \mathbb{P}^1$  is a finite morphism of degree  $\varphi^{-1}(\text{pt}) \times \phi^{-1}(\text{pt}) = D_i \cdot D_j = 1$ .  $\square$

The local version of Conjecture 1.1 was proved in [4, 18.22]:

**Theorem 2.3.** *Let  $(X, D = \sum d_i D_i)$  be a log pair which is log canonical at a point  $P \in \cap D_i$ . Assume that  $K_X$  and all  $D_i$ 's are  $\mathbb{Q}$ -Cartier at  $P$ . Then  $\sum d_i \leq \dim(X)$ . Moreover, if the equality holds, then  $(X \ni P, D)$  is an abelian quotient of a smooth point and  $(X, D)$  is not plt at  $P$ .*

Recall that for any plt pair  $(X, D)$  of dimension  $\leq 3$  there is a small birational contraction  $q: X^q \rightarrow X$  such that  $X^q$  is  $\mathbb{Q}$ -factorial and  $(X^q, D^q := q_*^{-1}D)$  is plt (see [4, 6.11.1], [4, 17.10]). Such  $q$  is called a  $\mathbb{Q}$ -factorialization of  $(X, D)$ .

Applying a  $\mathbb{Q}$ -factorialization in our situation and taking into account that  $\text{rk}(\text{Weil}(X)/\approx) = (\text{rk} \text{Weil}(X^q)/\approx) \geq \rho(X^q)$  we obtain that for Theorem 1.2 it is sufficient to prove the following

**Proposition 2.4.** *Let  $(X, D = \sum d_i D_i)$  be a three-dimensional projective plt pair such that  $K_X + D \equiv 0$  and  $X$  is  $\mathbb{Q}$ -factorial. Then*

$$(2.5) \quad \sum d_i \leq \rho(X) + 3.$$

*Moreover, if the equality holds, then for  $(X, \lfloor D \rfloor)$  there are only possibilities (i)–(iv) of Theorem 1.2.*

### 3. LEMMAS

In this section we prove several facts related to Conjecture 1.1.

**Lemma 3.1** (cf. [4, 18.24], [1]). *Let  $(X, D = \sum_{i=1}^r d_i D_i)$  be a projective  $n$ -dimensional log pair such that all  $D_i$ 's are  $\mathbb{Q}$ -Cartier,  $\rho(X) = 1$ ,  $(X, D)$  is plt and  $-(K_X + D)$  is nef. Then*

$$(3.2) \quad \sum d_i \leq n + 1.$$

*Moreover, if the equality holds, then  $X \simeq \mathbb{P}^n$  and  $D_1, \dots, D_r$  are hyperplanes.*

Note that in the two-dimensional case any plt pair is automatically  $\mathbb{Q}$ -factorial.

*Proof.* We will prove this lemma in the case when  $\lfloor D \rfloor = 0$  (i.e.  $K_X + D$  is klt). The case when  $\lfloor D \rfloor$  is non-trivial (and irreducible) can be treated in a similar way. The inequality (3.2) was proved in [4, 18.24], so we prove the second part of our lemma. Since  $-K_X$  is ample,  $\text{Pic}(X) \simeq \mathbb{Z}$  (see e.g. [8, 2.1.2]). Let  $H$  be an ample generator of  $\text{Pic}(X)$  and let  $D_i \equiv a_i H$ ,  $a_i \in \mathbb{Q}$ ,  $a_i \geq 0$ . Assume that  $a_i < a_j$  for  $i \neq j$  or  $K_X + D \not\equiv 0$ . For  $0 < \varepsilon \ll 1$ , consider

$$D^{(\varepsilon)} := \varepsilon D_i + D - \varepsilon D_j.$$

Then  $K_X + D^{(\varepsilon)}$  is again klt (because the klt property is an open condition) and  $-(K_X + D^{(\varepsilon)})$  is ample. Take  $N \in \mathbb{N}$  so that  $-N(K_X + D^{(\varepsilon)})$  is integral and very ample and let  $M \in |-N(K_X + D^{(\varepsilon)})|$  be a general member. By Bertini type theorem [3, Sect. 4],  $(X, D^{(\varepsilon)} + \frac{1}{N}M)$  is klt (and numerically trivial). Moreover, the sum of coefficients of  $D^{(\varepsilon)} + \frac{1}{N}M$  is equal to  $n + 1 + \frac{1}{N}$ . This contradicts (3.2). Hence  $K_X + D \equiv 0$  and  $D_i \equiv D_j$  for all  $i, j$ . Thus, for any pair  $D_i, D_j$  there exists  $n_{i,j} \in \mathbb{N}$  such that  $n_{i,j}(D_i - D_j) \sim 0$ . By taking repeated cyclic covers (which are étale in codimension one)  $\pi: X' \rightarrow \dots \rightarrow X$ , we obtain a new plt pair  $(X', D' = \sum_{i=1}^r d_i D'_i)$  [4, 20.4] such that  $D'_i \sim D'_j$ , where  $D'_i = \pi^* D_i$ . On this step, we do not assume that  $D'_i$  is irreducible. Then  $D'_1, \dots, D'_r$  generate a linear system  $\mathcal{L}$  of Weil divisors. If  $\text{Bs}(\mathcal{L})$  is not empty, then we pick a point  $P' \in D'_1 \cap \dots \cap D'_r$ . By construction,  $(X', D')$  is klt at  $P'$  and  $\sum_{i=1}^r d_i \geq n + 1$ , a contradiction with Theorem 2.3. Therefore  $\text{Bs}(\mathcal{L}) = \emptyset$ . In particular,  $D'_1, \dots, D'_r$  are ample Cartier

divisors and  $-K_{X'} \equiv D'$  is ample (i.e.  $X'$  is a log Fano variety). This also shows that the Fano index of  $X'$  is  $r(X') \geq \sum_{i=1}^r d_i \geq n+1$ . It is well known (see e.g. [8, 3.1.14]) that in this case we have  $r(X') = \sum_{i=1}^r d_i = n+1$ ,  $X' \simeq \mathbb{P}^n$  and  $D'_1, \dots, D'_r$  are hyperplanes. Since  $\pi: X' \rightarrow X$  is étale outside of  $\text{Sing}(X)$  and  $X'$  is smooth, the restriction  $X' \setminus \pi^{-1}(\text{Sing}(X)) \rightarrow X \setminus \text{Sing}(X)$  is the universal covering. This gives us that  $\pi: X' \rightarrow X$  is Galois. Hence  $X = \mathbb{P}^n/G$ , where  $G \subset PGL_{n+1}$  is a finite subgroup. Further, the group  $G$  does not permute  $D'_1, \dots, D'_r$ . Thus  $G$  has  $r > n+1$  invariant hyperplanes  $D'_1, \dots, D'_r$  in  $\mathbb{P}^n$ . Finally, the lemma follows by the following simple fact which can be proved by induction on  $n$ .  $\square$

**Sublemma.** *Let  $G \subset PGL_{n+1}$  be a finite subgroup acting on  $\mathbb{P}^n$  free in codimension one. Assume that there are  $n+2$  invariant hyperplanes  $H_1, \dots, H_{n+2} \subset \mathbb{P}^n$ . Then  $G = \{1\}$ .*

**Lemma 3.3.** *Let  $\varphi: X \rightarrow Z \ni o$  be a three-dimensional flipping contraction and let  $D = \sum d_i D_i$  be a boundary on  $X$  such that  $(X, D)$  is plt,  $\rho(X/Z) = 1$ ,  $-(K_X + D)$  is  $\varphi$ -nef and all  $D_i$ 's are  $\varphi$ -ample. Assume that  $X$  is  $\mathbb{Q}$ -factorial. Then  $\sum d_i < 2$ .*

*Proof.* Let  $\chi: X \xrightarrow{\varphi} Z \xleftarrow{\varphi^+} X^+$  be the flip with respect to  $K_X$  and let  $D^+ = \sum d_i D_i^+$  be the proper transform of  $D$ . Then all  $D_i^+$ 's are anti-ample over  $Z$ . Hence  $\varphi^{+-1}(o)$  is contained in  $\cap D_i^+$ . Consider a general hyperplane section  $H \subset X^+$ . Then  $(H, D|_H)$  is plt [3, Sect. 4]. Applying Theorem 2.3 to  $H$  we obtain  $\sum d_i < 2$ .  $\square$

**Lemma 3.4.** *Let  $\varphi: X \rightarrow Z$  be a contraction from a projective  $\mathbb{Q}$ -factorial threefold onto a surface such that  $\rho(X/Z) = 1$ . Let  $D = \sum d_i D_i$  be a boundary on  $X$  such that  $(X, D)$  is lc,  $(X, D - \lfloor D \rfloor)$  is klt and  $-(K_X + D)$  is nef. Assume that  $\lfloor D \rfloor$  has a component  $S$  which is generically section of  $\varphi$ . Then  $\sum_i d_i \leq \rho(X) + 3$ . Moreover, if the equality holds and  $(X, D)$  is plt, then  $X$  is smooth,  $\varphi$  is a  $\mathbb{P}^1$ -bundle,  $\varphi|_S: S \rightarrow Z$  is an isomorphism and  $Z \simeq \mathbb{P}^2$  or  $Z \simeq \mathbb{P}^1 \times \mathbb{P}^1$ .*

*Proof.* Assume that  $\sum_i d_i \geq \rho(X) + 3$ . Since  $-K_X$  is  $\varphi$ -ample, a general fiber  $\ell$  of  $\varphi$  is isomorphic to  $\mathbb{P}^1$ . We have

$$(3.5) \quad 2 = -K_X \cdot \ell = D^{\text{hor}} \cdot \ell \geq \sum_{\text{hor}} d_i, \quad \sum_{\text{ver}} d_i \geq \rho(X) + 1 = \rho(Z) + 2.$$

Let  $\mu := \varphi|_S$ . Write  $\text{Diff}_S(D - S) = \sum_i \beta_i \Theta_i$ . Then

$$(3.6) \quad \beta_i = 1 - \frac{1}{m_i} + \frac{1}{m_i} \sum_{j \in \mathfrak{M}_i} d_j k_{i,j},$$

where  $m_i \in \mathbb{N} \cup \{\infty\}$ ,  $k_{i,j} \in \mathbb{N}$  and the sum runs through the set  $\mathfrak{M}_i$  of all components  $D_j$  containing  $\Theta_i$  (see [11, 3.10]). Here  $m_i = \infty$  when  $(X, D)$  is not plt along  $\Theta_i$ . It is easy to see that  $\beta_i \geq \sum_{j \in \mathfrak{M}_i} d_j$ . Put  $\Xi := \mu_* \text{Diff}_S(D - S)$  and

let  $\Xi = \sum \gamma_i \Xi_i$ . Since  $-(K_S + \Theta)$  is nef,  $(Z, \Xi)$  is lc [4, 2.28]. For any component  $D_i$  of  $D^{\text{ver}}$  we have at least one component  $\Theta_j \subset D_i \cap S$  such that  $\mu(\Theta_j) \neq \text{pt}$ . This yields

$$\sum_i \gamma_i \geq \sum_{\mu(\Theta_i) \neq \text{pt}} \beta_i \geq \sum_{\text{ver}} d_j \geq \rho(Z) + 2.$$

Applying Proposition 2.1 to  $(Z, \Xi)$ , we obtain equalities

$$(3.7) \quad \sum_i \gamma_i = \sum_{\mu(\Theta_i) \neq \text{pt}} \beta_i = \sum_{\text{ver}} d_j = \rho(Z) + 2.$$

Hence  $\sum_{\text{hor}} d_i = 2$  and  $\sum_i d_i = \rho(X) + 3$ . This shows the first part of the lemma.

Now assume that  $(X, D)$  is plt. By Adjunction [4, 17.6],  $(S, \text{Diff}_S(D - S))$  is klt and so is  $(Z, \Xi)$ . Again, by Proposition 2.1 we have either  $Z \simeq \mathbb{P}^2$  or  $Z \simeq \mathbb{P}^1 \times \mathbb{P}^1$ . There exists a *standard form* of  $\varphi$  (see [10]), i. e. the commutative diagram

$$\begin{array}{ccc} \tilde{X} & \dashrightarrow & X \\ \downarrow & & \downarrow \\ \tilde{Z} & \xrightarrow{\sigma} & Z \end{array}$$

where  $\sigma: \tilde{Z} \rightarrow Z$  is a birational morphism of smooth surfaces,  $\tilde{X} \dashrightarrow X$  is a birational map and  $\tilde{\varphi}: \tilde{X} \rightarrow \tilde{Z}$  is a *standard* conic bundle (in particular,  $\tilde{X}$  is smooth and  $\rho(\tilde{X}/\tilde{Z}) = 1$ ). Take the proper transform  $\tilde{S}$  of  $S$  on  $\tilde{X}$ . For a general fiber  $\tilde{\ell}$  of  $\tilde{\varphi}$  we have  $\tilde{S} \cdot \tilde{\ell} = 1$ . Since  $\rho(\tilde{X}/\tilde{Z}) = 1$ ,  $\tilde{S}$  is  $\tilde{\varphi}$ -ample. It gives us that each fiber of  $\tilde{\varphi}$  is reduced and irreducible, i. e. the morphism  $\tilde{\varphi}$  is smooth. By [7], there exists a standard conic bundle  $\hat{\varphi}: \hat{X} \rightarrow Z$  and a birational map  $\hat{X} \dashrightarrow X$  over  $Z$ . This map induces an isomorphism  $(\hat{X}/\hat{\varphi}^{-1}(\mathfrak{M})) \simeq (X/\varphi^{-1}(\mathfrak{M}))$ , where  $\mathfrak{M} \subset Z$  is a finite number of points. Since both  $\varphi, \hat{\varphi}$  are projective and  $\rho(X/Z) = \rho(\hat{X}/Z) = 1$ , we have  $\hat{X} \simeq X$ . But then  $\varphi: X \rightarrow Z$  is smooth, i.e.  $\varphi$  is a  $\mathbb{P}^1$ -bundle.

Now we claim that  $\mu$  is an isomorphism. Indeed, otherwise  $S$  contains a fiber, say  $\ell_0$ . Then  $S$  intersects all irreducible components of  $D^{\text{hor}} - S$ . If some component  $D_k$  of  $D^{\text{hor}} - S$  does not contain  $\ell_0$ , then  $\varphi(S \cap D_k)$  is a component of  $\Xi$ . By (3.7) we have

$$\rho(Z) + 2 = \sum_i \gamma_i = \sum_{\mu(\Theta_i) \neq \text{pt}} \beta_i \geq d_k + \sum_{\text{ver}} d_j > \rho(Z) + 2,$$

which is impossible. Therefore all components of  $D^{\text{hor}}$  contain  $\ell_0$ . Taking a general hyperplane section as in the proof of Lemma 3.3, we derive a contradiction.  $\square$

**Corollary 3.7.1.**  *$S$  does not intersect  $\text{Supp}(D^{\text{hor}} - S)$  and all components of  $D^{\text{hor}} - S$  are sections of  $\varphi$ .*

**Lemma 3.8.** *Let  $\varphi: X \rightarrow Z$  be a contraction from a  $\mathbb{Q}$ -factorial threefold onto a curve and let  $D = \sum d_i D_i$  be a boundary on  $X$  such that  $(X, D)$  is lc,  $(X, D - \lfloor D \rfloor)$  is klt. Let  $F$  be a general fiber. Assume that  $-(K_X + D)$  is  $\varphi$ -nef and  $\rho(X/Z) = 1$ . Then  $\sum_{\text{hor}} d_i \leq 3$ . Moreover, if the equality holds and  $(X, D)$  is plt, then  $F \simeq \mathbb{P}^2$  and for any component  $D_i$  of  $D^{\text{hor}}$  the scheme-theoretic restriction  $D_i|_F$  is a line.*

*Proof.* Put  $\Delta := D|_F$ . Then  $(F, \Delta)$  is lc,  $(F, \Delta - \lfloor \Delta \rfloor)$  is klt (see [3, Sect. 4]) and  $-(K_F + \Delta)$  is nef. Moreover, if  $(X, D)$  is plt, then so is  $(F, \Delta)$ . Write  $\Delta = \sum \delta_i \Delta_i$ , where all  $\Delta_i$ 's are irreducible curves on  $F$ . Clearly  $D^{\text{ver}}|_F = 0$  and  $\sum \delta_i \geq \sum_{\text{hor}} d_i$ . If  $\rho(F) = 1$ , then the assertion of 3.8 follows by Proposition 2.1. Assume that  $\rho(F) > 1$ . Let  $C$  be an extremal  $K_F$ -negative curve on  $F$  (note that  $K_F$  is not nef). Then  $C$  intersects all components of  $\Delta$  (because  $\rho(X/Z) = 1$ ). Let  $v: F \rightarrow F'$  be the contraction of  $C$ . If  $F'$  is a curve, then we take  $C$  to be a general fiber of  $v$ . By Adjunction,  $2 = -\deg K_C \geq \deg \Delta|_C$ . This gives us  $2 \geq \sum \delta_i \geq \sum_{\text{hor}} d_i$ . If  $v$  is birational, then  $(F', v(\Delta))$  is lc and all components of  $v(\Delta)$  pass through the point  $v(C)$ . By Theorem 2.3, the sum of coefficients of  $v(\Delta)$  is  $\leq 2$ . Hence  $\sum_{\text{hor}} d_i \leq \sum \delta_i \leq 3$ . If  $(F, \Delta)$  is klt, then so is  $(F', v(\Delta))$  and the inequality above is strict. Finally, if  $(F, \Delta)$  is plt and  $\lfloor \Delta \rfloor \neq 0$ , then we take  $C$  to be  $(K_F + \Delta - \lfloor \Delta \rfloor)$ -negative extremal curve. Then  $C$  is not a component of  $\lfloor \Delta \rfloor$ . By [3, 3.10],  $(F', v(\Delta))$  is plt. Again, by Theorem 2.3 the sum of coefficients of  $v(\Delta)$  is strictly less than 2. So,  $\sum_{\text{hor}} d_i \leq \sum \delta_i < 3$ . This proves Lemma 3.8.  $\square$

**Corollary 3.8.1.** *Notation as in Lemma 3.8. Assume additionally that  $X$  is projective,  $-(K_X + D)$  is nef (not only over  $Z$ ),  $\sum_{\text{hor}} d_i = 3$ ,  $\sum_{\text{ver}} d_i = 2$  and  $(X, D)$  is plt. If  $\lfloor D^{\text{hor}} \rfloor \neq \emptyset$ , then  $\lfloor D^{\text{hor}} \rfloor = \lfloor D \rfloor \simeq \mathbb{P}^1 \times \mathbb{P}^1$  and  $X$  is smooth along  $\lfloor D \rfloor$ . In particular,  $X$  has at most isolated singularities.*

*Proof.* Put  $S := \lfloor D \rfloor$ . By [4, 17.5],  $S$  is normal. Since  $\rho(X/Z) = 1$ ,  $S$  is irreducible and all components of  $D - S$  meet  $S$ . Let  $\text{Diff}_S(D - S) = \sum \beta_i \Theta_i$ . Clearly,  $-(K_S + \text{Diff}_S(D - S))$  is nef. By [4, 17.6],  $(S, \Theta)$  is klt. As in the proof of Lemma 3.4 we see  $\sum \beta_i \geq \sum d_i - 1 \geq 4$ . If  $\rho(S) = 2$ , then equalities  $\sum \beta_i = \sum d_i - 1 = 4$  and Proposition 2.1 give us the assertion. Assume that  $\rho(S) > 2$ . Then some fiber of  $\varphi|_S: S \rightarrow Z$  is not irreducible. Let  $\Gamma$  be its irreducible component and let  $v: S \rightarrow S'$  be the contraction of  $\Gamma$ . Taking into account that  $\Gamma$  intersects all components of  $D^{\text{hor}}$ . As in Lemma 3.4 we get a contradiction.  $\square$

**Lemma 3.9** (cf. [11, 6.9]). *Let  $\varphi: X \rightarrow Z \ni o$  be a  $K_X$ -negative contraction from a  $\mathbb{Q}$ -factorial variety  $X$  such that  $\rho(X/Z) = 1$  and every fiber has dimension one. Let  $D$  be a boundary on  $X$  such that  $(X, D - \lfloor D \rfloor)$  is klt and  $K_X + D$  is  $\varphi$ -numerically trivial. Assume that  $\lfloor D \rfloor$  is disconnected near  $\varphi^{-1}(o)$ . Then  $K_X + D$  is plt near  $\varphi^{-1}(o)$ .*

*Proof.* Regard  $\varphi: X \rightarrow Z \ni o$  as a germ near  $\varphi^{-1}(o)$ . Put  $S := \lfloor D \rfloor$ . Clearly, for a general fiber  $\ell$  of  $\varphi$  we have  $-K_X \cdot \ell = D \cdot \ell = 2$ . If  $S'$  is an irreducible component

of  $S$  such that  $S' \cdot \ell = 0$ , then  $S' = \varphi^{-1}(C)$  for a curve  $C \subset Z$ . In this case,  $S'$  contains  $\varphi^{-1}(o)$  and  $S$  is connected near  $\varphi^{-1}(o)$ . Therefore  $S$  has exactly two connected components  $S_1, S_2$ , they are irreducible and  $S_1 \cdot \ell = S_2 \cdot \ell = 1$ . Then  $S_i$ ,  $i = 1, 2$  meets all components of  $\varphi^{-1}(o)$ . Hence  $S_i \cap \varphi^{-1}(o)$  is 0-dimensional. Since  $Z$  is normal and  $\varphi|_{S_i}: S_i \rightarrow Z$  is birational,  $S_i \simeq Z$  and  $S_i \cap \varphi^{-1}(o)$  is a single point. In particular,  $\varphi^{-1}(o)$  is irreducible. Clearly,  $LCS(X, D) \subset S = S_1 \cup S_2$ . Assume that  $(X, D)$  is not plt. Then there is a divisor  $E$  of the function field  $K(X)$  with discrepancy  $a(E, D) < -1$ . Let  $V \subset X$  be its center. Then  $V \subset S$  and we may assume that  $V \subset S_1$  (and  $V \neq S_1$ ). Let  $L \subset Z$  be any effective prime divisor containing  $\varphi(V)$  and let  $F := \varphi^{-1}(L)$ . Clearly,  $(X, D + F)$  is not lc near  $V$ . For sufficiently small positive  $\varepsilon$  the log pair  $(X, D + F - \varepsilon S_1)$  is not lc near  $V$  and not klt near  $S_2$ . This contradicts Connectedness Lemma [4, 17.4]  $\square$

#### 4. PROOF OF THEOREM 1.2

In this section we prove Proposition 2.4.

**4.1. Inductive hypothesis.** Notation and assumption as in Proposition 2.4. Our proof is by induction on  $\rho(X)$ . In the case  $\rho(X) = 1$ , the assertion is a consequence of Lemma 3.1. To prove 2.4 for  $\rho(X) \geq 2$  we fix  $\rho \in \mathbb{N}$ ,  $\rho > 1$ . Assume that inequality (2.5) holds if  $\rho(X) < \rho$  and for  $\rho(X) = \rho$  we have

$$(4.2) \quad \sum d_i - \rho(X) - 3 \geq 0.$$

**4.3.** If  $(X, D)$  is klt, then we run  $K_X$ -MMP. On each step  $K \equiv -D$  cannot be nef. Obviously, all steps preserve our assumptions (see [4, 2.28]) and the left hand side of (4.2) does not decrease. Moreover, by our assumptions we have no divisorial contractions on  $X$  (because after any divisorial contraction the left hand side of (4.2) decreases). Therefore after a number of flips, we obtain a fiber type contraction  $\varphi: X \rightarrow Z$ . Since  $\rho(X) = \rho \geq 2$ ,  $\dim(Z) = 1$  or  $2$ . Note that all varieties from Theorem 1.2 have no small contractions. Thus, it is sufficient to prove Proposition 2.4 on our new model  $(X, D)$ .

This procedure does not work if  $(X, D)$  is not klt. The difference is that contractions of components of  $D$  do not contradict the inductive hypothesis. If  $(X, D)$  is not klt, then we run  $(K_X + D - \lfloor D \rfloor)$ -MMP. Note that  $\lfloor D \rfloor$  is normal and irreducible [4, 17.5]. For every extremal ray  $R$  we have  $\lfloor D \rfloor \cdot R > 0$ , so we cannot contract an irreducible component of  $\lfloor D \rfloor$ . Therefore after every divisorial contraction  $\sum d_i - \rho(X)$  decrease, a contradiction with our assumption. Thus, all steps of the MMP are flips. By [4, 2.28], they preserve the plt property of  $K + D$ . At the end we get a fiber type contraction  $\varphi: X \rightarrow Z$ , where  $\dim(Z) < 3$  and  $\lfloor D \rfloor$  is  $\varphi$ -ample (i.e.  $\lfloor D^{\text{hor}} \rfloor \neq 0$ ). Since  $\lfloor D^{\text{hor}} \rfloor$  has a component which intersects all components of  $D^{\text{ver}}$ ,  $\lfloor D^{\text{ver}} \rfloor = 0$ .



4.4. **Case:**  $\dim(Z) = 1$ . Then  $\rho(X) = 2$ . By Lemma 3.8 and our assumption (4.2), we have  $\sum_{\text{hor}} d_i \leq 3$  and  $\sum_{\text{ver}} d_i \geq 2$ . In particular,  $D^{\text{ver}} \neq 0$ . Components of  $D^{\text{ver}}$  are fibers of  $\varphi$ , so they are numerically proportional. Clearly, the log divisor  $K_X + D^{\text{hor}} \equiv -D^{\text{ver}}$  is not nef and curves in fibers of  $\varphi$  are trivial with respect to it. Let  $Q$  be the extremal  $(K_X + D^{\text{hor}})$ -negative ray of  $\overline{NE}(X) \subset \mathbb{R}^2$  and let  $\phi: X \rightarrow W$  be its contraction. It follows by Lemma 3.3 that  $\phi$  cannot be a flipping contraction. Let  $\ell$  be a general curve such that  $\phi(\ell) = \text{pt.}$  Then  $\ell$  dominates  $Z$  and  $\ell \simeq \mathbb{P}^1$ . Hence  $Z \simeq \mathbb{P}^1$ .

4.4.1. *Subcase*  $[D] = 0$ . We will prove that  $X \simeq \mathbb{P}^2 \times \mathbb{P}^1$ . By our inductive hypothesis,  $\phi$  cannot be divisorial. Therefore  $\dim(W) = 2$ . Further,  $D^{\text{ver}} \cdot \ell \leq D \cdot \ell = -K_X \cdot \ell = 2$ . Since  $\ell$  intersects all components of  $D^{\text{ver}}$ ,  $\sum_{\text{ver}} d_i \leq 2$ . This yields  $\sum_{\text{ver}} d_i = 2$  and  $\sum_{\text{hor}} d_i = 3$ . In particular, this proves inequality (2.5). Moreover,  $\ell \cdot D^{\text{ver}} = 2$ ,  $\ell \cdot D^{\text{hor}} = 0$  and for any component  $D_i$  of  $D^{\text{ver}}$  we have  $D_i \cdot \ell = 1$ . Fix two components of  $D^{\text{ver}}$ , say  $D_0$  and  $D_1$ . Then  $K_X + D_0 + D_1 + D^{\text{hor}} \equiv K_X + D \equiv 0$ , so  $(X, D_0 + D_1 + D^{\text{hor}})$  is plt by Lemma 3.9. Applying Lemma 3.4 we obtain  $D_0 \simeq D_1 \simeq Z \simeq \mathbb{P}^2$ ,  $X$  is smooth and  $\phi$  is a  $\mathbb{P}^1$ -bundle. By [6, 3.5],  $\varphi$  is a  $\mathbb{P}^2$ -bundle. We have a finite morphism  $\varphi \times \phi: X \rightarrow Z \times W = \mathbb{P}^1 \times \mathbb{P}^2$ . Clearly,  $\deg(\varphi \times \phi) = \varphi^{-1}(\text{pt}) \cdot \ell = 1$ . Hence  $\varphi \times \phi$  is an isomorphism.

4.4.2. *Subcase:*  $[D] \neq 0$ . Since  $\rho(X/Z) = 1$ ,  $[D]$  is irreducible. Put  $S := [D]$ . Let  $F$  be a general fiber of  $\varphi$ . By construction,  $-K_X$  is  $\varphi$ -ample. First, assume that  $\dim(W) = 2$ . Then  $D^{\text{ver}} \cdot \ell \leq D \cdot \ell = -K_X \cdot \ell = 2$ . Since  $\ell$  intersects all components of  $D^{\text{ver}}$ ,  $\sum_{\text{ver}} d_i \leq 2$ . This yields  $\sum_{\text{ver}} d_i = 2$ ,  $\sum_{\text{hor}} d_i = 3$  and  $\sum d_i = 5$ . Moreover,  $D^{\text{hor}} \cdot \ell = 0$ . By Lemma 3.8,  $F \simeq \mathbb{P}^2$ ,  $X$  is smooth along  $F$  and for any component  $D_i$  of  $D^{\text{hor}}$  the scheme-theoretic restriction  $D_i|_F$  is a line. Hence components of  $D^{\text{hor}}$  are numerically equivalent. Let  $D_1$  be a component of  $D^{\text{hor}} - S$ . Consider the new boundary  $D' := D + \varepsilon D_0 - \varepsilon S$ . If  $0 < \varepsilon \ll 1$ , then  $(X, D')$  is klt and  $K_X + D' \equiv 0$ . Applying Case 4.4.1 we get  $W \simeq \mathbb{P}^2$  and  $X \simeq \mathbb{P}^2 \times \mathbb{P}^1$ .

Now assume that  $\phi$  is divisorial. By the inductive hypothesis,  $\phi$  contract  $S$ . Since the contraction is extremal,  $\phi(S)$  is a curve (otherwise curves  $S \cap \varphi^{-1}(\text{pt})$  is contracted by  $\phi$  and  $\varphi$ ). All components of  $\phi(D^{\text{ver}})$  pass through  $\phi(S)$ . By taking a general hyperplane section as in the proof of Lemma 3.3, we obtain  $\sum_{\text{ver}} d_i \leq 2$ . By Corollary 3.8.1, we obtain that  $S \simeq \mathbb{P}^1 \times \mathbb{P}^1$ ,  $X$  has only isolated singularities and  $X$  is smooth along  $S$ . By Lemma 3.8,  $F \simeq \mathbb{P}^2$  and  $X$  is smooth along  $F$ . The curve  $F \cap S$  is ample on  $F$ , so it is connected and smooth by the Bertini theorem. Therefore  $F \cap S$  is a generator of  $S = \mathbb{P}^1 \times \mathbb{P}^1$ . Since  $\varphi|_S$  is flat, he same holds for arbitrary fiber  $F_0$ . Hence all fibers of  $\varphi$  are numerically equivalent and any fiber  $F_0$  contains an ample smooth rational curve. Moreover, this also means  $F_0$  is not multiple. Thus it is a normal surface. Now as in Case 4.4.1,  $K_X + F_0 + F_1 + D^{\text{hor}} \equiv 0$  and by Lemma 3.9,  $(X, F_0 + F_1 + D^{\text{hor}})$  is plt for any fibers  $F_0, F_1$ . By Adjunction,  $(F_0, D^{\text{hor}}|_{F_0})$  is klt. Clearly,  $K_{F_0} \equiv K_X|_{F_0}$  and  $S|_{F_0}$

are numerically proportional. Hence  $F_0$  is a log del Pezzo surface of Fano index  $> 1$ . Since  $\varphi$  is flat,  $(K_{F_0})^2 = (K_F)^2 = 9$ . Therefore,  $F_0 \simeq \mathbb{P}^2$  and  $X$  is smooth. By [6, 3.5],  $\varphi$  is a  $\mathbb{P}^2$ -bundle so,  $X \simeq \mathbb{P}_{\mathbb{P}^1}(\mathcal{E})$ , where  $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(b)$ ,  $0 \leq a \leq b$ . The Grothendieck tautological bundle  $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$  is generated by global sections and not ample. Therefore  $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$  gives us a supporting function for the extremal ray  $Q$ . Since  $\phi$  is birational,  $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)^3 = a + b > 0$ . Finally,  $X$  contains  $S = \mathbb{P}^1 \times \mathbb{P}^1$ . Hence  $a = 0$ . This proves Proposition 2.4 in the case when  $Z$  is a curve.

**4.5. Case:**  $\dim(Z) = 2$ . Note that  $Z$  has only log terminal singularities (see e. g. [4, 15.11]). Since  $-K_X$  is  $\varphi$ -ample, a general fiber  $\ell$  of  $\varphi$  is  $\mathbb{P}^1$ . Hence  $2 = -K_X \cdot \ell = D \cdot \ell = D^{\text{hor}} \cdot \ell \geq \sum_{\text{hor}} d_i$ . By our assumption,  $\sum_{\text{ver}} d_i \geq \rho(X) + 1$ . If  $(X, D)$  is not plt, then  $\lfloor D \rfloor$  is  $\varphi$ -ample. Clearly,  $\lfloor D^{\text{ver}} \rfloor = 0$ .

**Claim 4.5.1.** *Notation as above. Then  $K_Z$  is not nef.*

*Proof.* Run  $(K_X + D^{\text{ver}})$ -MMP. After a number of flips we get either a divisorial contraction (of the proper transform of a component of  $\lfloor D \rfloor$ ), or a fiber type contraction. In both cases  $Z$  is dominated by a family of rational curves [2, 5-1-4, 5-1-8]. Therefore  $K_Z$  is not nef by [5].  $\square$

**Claim 4.5.2.** *Notation as in 4.5. Then  $Z$  contains no contractible curves. In particular,  $\rho(Z) \leq 2$ .*

*Proof.* Assume the converse. i. e. there is an irreducible curve  $\Gamma \subset Z$  and a birational contraction  $\mu: Z \rightarrow Z''$  such that  $\mu(\Gamma) = \text{pt}$  and  $\rho(Z/Z'') = 1$ . Denote  $F := \varphi^{-1}(\Gamma)$ . Since  $F \not\subset \lfloor D \rfloor$ ,  $(X, D + \varepsilon F)$  is plt for  $0 < \varepsilon \ll 1$  [4, 2.17]. First we assume that  $\Gamma$  is contractible: Run  $(K + D + \varepsilon F)$ -MMP over  $Z''$ . By our inductive hypothesis, there are no divisorial contractions (because such a contraction must contract  $F$ ). At the end we cannot get a fiber type contraction (because  $K + D + \varepsilon F \equiv \varepsilon F$  cannot be anti-ample over a lower-dimensional variety). Thus after a number of flips  $X \dashrightarrow X'$ , we get a model  $X'$  over  $Z''$  such that  $K_{X'} + D' + \varepsilon F' \equiv \varepsilon F'$  is nef over  $Z''$ , where  $D'$  and  $F'$  are proper transforms of  $D$  and  $F$ , respectively. Then  $F' \neq 0$  (because  $F \neq 0$ ). Let  $\ell'$  be the proper transform of a general fiber of  $\varphi$ . Since  $F'$  is nef over  $Z''$ ,  $F' \cdot \ell' = 0$  and  $\rho(X'/Z'') = 2$ , we obtain that  $\ell'$  generates an extremal ray of  $\overline{NE}(X'/Z'')$ . Let  $\varphi': X' \rightarrow Z'$  be its contraction and let  $\mu': Z' \rightarrow Z''$  be the natural map. Then  $\dim(Z') = 2$ ,  $\Gamma' := \varphi'(F')$  is a curve and  $\mu'(\Gamma') = \mu(\varphi(F)) = \mu(\Gamma) = \text{pt}$ . Therefore  $(\Gamma')^2 < 0$ . On the other hand,  $(\Gamma')^2 \geq 0$ , which is a contradiction. Indeed, Let  $C' \subset F'$  be any curve such that  $\varphi'(C') = \Gamma'$ . Then  $C' \cdot F' \geq 0$ . By the projection formula,  $(\Gamma')^2 \geq 0$ .

Since  $K_Z$  is not nef, there is an extremal contraction  $\psi: Z \rightarrow V$ . By the above it is not birational. Therefore  $\dim(V) = 1$  and  $\rho(Z) = 2$ .  $\square$

**Corollary 4.5.3.** *Notation as in 4.5. One of the following holds:*

- (i)  $\rho(Z) = 1$  and  $-K_Z$  is ample;
- (ii)  $\rho(Z) = 2$  and there is a  $K_Z$ -negative extremal contraction  $\psi: Z \rightarrow V$  onto a curve.

4.5.4. *Subcase  $[D] = 0$ .* Let  $D_i$  be a component of  $D^{\text{ver}}$ . Run  $(K + D - d_i D_i)$ -MMP:

$$\chi_{(i)}: X \dashrightarrow X^{(i)}.$$

As above we get a fiber type contraction  $\varphi_{(i)}: X^{(i)} \rightarrow Z^{(i)}$ . Notations  $D^{\text{ver}}$  and  $D^{\text{hor}}$  will be fixed with respect to our original  $\varphi$ . If  $\dim(Z^{(i)}) = 1$ , then replacing  $X$  with  $X^{(i)}$  we get the case  $\dim(Z) = 1$  above. Thus we can assume that  $\dim(Z^{(i)}) = 2$  for any chose of  $D_i$ . Let  $\ell^{(i)} \subset X^{(i)}$  be a general fiber of  $\varphi_{(i)}$  and let  $L^{(i)} \subset X$  be its proper transform. Clearly,  $\chi_{(i)}$  is an isomorphism along  $L^{(i)}$ . Hence  $-K_X \cdot L^{(i)} = 2$ ,  $L^{(i)}$  is nef and  $D_i \cdot L^{(i)} > 0$ . For  $i = 1, \dots, r$  we get rational curves  $L^{(1)}, \dots, L^{(r)}$ . We shift indexing so that  $X = X^{(0)}$  and put  $Z = Z^{(0)}$  and  $\varphi = \varphi_{(0)}$ .

Up to permutations we can take  $L^{(0)}, \dots, L^{(s)}$ ,  $s+1 \leq r$  to be linearly independent in  $N_1(X)$ . Then for any  $D_i$  there exists  $L^{(j)}$  such that  $D_i \cdot L^{(j)} > 0$ . Thus we have

$$\begin{aligned} 2(s+1) &= -K_X \cdot \sum_{j=0}^s L^{(j)} = D \cdot \sum_{j=0}^s L^{(j)} = \\ &= \sum_{i=1}^r d_i \left( D_i \cdot \sum_{j=0}^s L^{(j)} \right) \geq \sum_{i=1}^r d_i \geq \rho(X) + 3. \end{aligned}$$

Since  $3 \geq \rho(Z) + 1 = \rho(X) \geq s+1$ , this yields  $\rho(X) = s+1 = 3$ . Thus,

$$(4.6) \quad D_i \cdot \sum_{j=0}^2 L^{(j)} = 1$$

holds for all  $i$ . Moreover,  $L^{(0)}, L^{(1)}, L^{(2)}$  generate  $N_1(X)$  and components of  $D$  generate  $N^1(X)$ .

Taking into account that  $2 = -K_X \cdot L^{(j)} = D \cdot L^{(j)}$ , we decompose  $D$  into the sum  $D = D^{(0)} + D^{(1)} + D^{(2)}$  of effective  $\mathbb{Q}$ -divisors without common components so that

$$(4.7) \quad D^{(i)} \cdot L^{(j)} = \begin{cases} 0 & \text{if } i \neq j \\ 2 & \text{otherwise} \end{cases}$$

Then  $D^{(i)} = \varphi^* \Delta^{(i)}$  for  $i = 1, 2$ , where  $\Delta^{(1)}, \Delta^{(2)}$  are effective  $\mathbb{Q}$ -divisors on  $Z$ . Put  $C^{(i)} := \varphi(L^{(i)})$ ,  $i = 1, 2$ . Since families  $L^{(j)}$  are dense on  $X$ ,  $C^{(j)}$  are nef and

$\neq 0$ . By the projection formula,

$$\Delta^{(i)} \cdot C^{(j)} \quad \begin{cases} = 0 & \text{if } 1 \leq i \neq j \leq 2 \\ > 0 & \text{if } 1 \leq i = j \leq 2 \end{cases}$$

Hence  $\Delta^{(1)}$  and  $\Delta^{(2)}$  generate extremal rays of  $\overline{NE}(Z) \subset \mathbb{R}^2$ . By (4.7), these  $\mathbb{Q}$ -divisors have more than one component, so they are nef and  $(\Delta^{(1)})^2 = (\Delta^{(2)})^2 = 0$ . This gives us that  $C^{(1)}$  and  $C^{(2)}$  also generate extremal rays. Therefore  $C^{(i)}$  and  $\Delta^{(j)}$  are numerically proportional whenever  $i \neq j$  and  $(C^{(1)})^2 = (C^{(2)})^2 = 0$ . In particular,  $C^{(i)}$ ,  $i = 1, 2$  generate an one-dimensional base point free linear system which defines a contraction  $Z \rightarrow \mathbb{P}^1$ . This also shows that  $D^{(i)} = \varphi^* \Delta^{(i)}$ ,  $i = 1, 2$  are nef on  $X$ .

Now we claim that  $D^{(0)}$  is nef. Assume the opposite. Then for small  $\varepsilon > 0$ ,  $(X, D + \varepsilon D^{(0)})$  is klt [4, 2.17]. There is a  $(K_X + D + \varepsilon D^{(0)})$ -negative extremal ray, say  $R$ . By our inductive hypothesis, the contraction of  $R$  must be of flipping type. Since  $\Delta^{(1)}, \Delta^{(2)}$  generate  $N^1(Z)$ , we have  $D^{(i)} \cdot R > 0$  for  $i = 1$  or  $2$ . By (4.7),  $\sum_j^{(i)} d_j = 2$ , where  $\sum_j^{(i)}$  runs through all components  $D_j$  of  $D^{(i)}$ . Since components of  $D^{(i)} = \varphi^*(\Delta^{(i)})$  are numerically proportional, this contradicts Lemma 3.3. Therefore  $D^{(i)}$  are nef for  $i = 0, 1, 2$ .

We claim that  $L^{(i)}$ ,  $i = 0, 1, 2$  generate  $\overline{NE}(X)$ . Indeed, let  $z \in \overline{NE}(X)$  be any element. Then  $z \equiv \sum \alpha_i [L^{(i)}]$  for  $\alpha_i \in \mathbb{R}$ . By (4.7),  $0 \leq D^{(j)} \cdot z = \alpha_j$ . This shows that  $L^{(i)}$  generate  $\overline{NE}(X)$ . From (4.6) we see that components of  $D^{(0)}$  are numerically equivalent.

Fix two components  $D'$  and  $D''$  of  $D^{(0)}$ . Then  $K_X + D' + D'' + D^{(1)} + D^{(2)} \equiv 0$ . By Lemma 3.9,  $(X, D' + D'' + D^{(1)} + D^{(2)})$  is plt. Lemma 3.4 implies that  $D' \simeq D'' \simeq S \simeq \mathbb{P}^1 \times \mathbb{P}^1$ ,  $X$  is smooth and  $\varphi$  is a  $\mathbb{P}^1$ -bundle. As in the case  $\dim(Z) = 1$ , we have  $X \simeq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ .

**4.7.5. Subcase  $[D] \neq 0$ .** Let  $S$  be a component of  $[D]$ . Clearly,  $S \cdot \ell \leq 2$ . If  $S$  is generically a section of  $\varphi$ , then by Lemma 3.4,  $X$  is smooth,  $\varphi$  is a  $\mathbb{P}^1$ -bundle and  $S \simeq \mathbb{P}^2$  or  $\mathbb{P}^1 \times \mathbb{P}^1$ . Therefore  $X \simeq \mathbb{P}(\mathcal{E})$ , where  $\mathcal{E}$  is a rank two vector bundle on  $Z$ . Since  $\varphi$  has disjoint sections,  $\mathcal{E}$  is decomposable. So we may assume that  $\mathcal{E} = \mathcal{O}_Z + \mathcal{L}$ , where  $\mathcal{L}$  is a line bundle. By the projection formula, all components of  $D^{\text{ver}}$  are nef. Let  $R$  be a  $(K_X + D^{\text{hor}})$ -negative extremal curve and let  $\phi: X \rightarrow W$  be its contraction. Assume that  $\phi$  is of flipping type. By [6],  $K_X \cdot R \geq 0$ . Hence  $D^{\text{hor}} \cdot R < 0$ , so  $R$  is contained in a section of  $\varphi$ . But all curves on  $\mathbb{P}^2$  and  $\mathbb{P}^1 \times \mathbb{P}^1$  are movable, a contradiction. If  $\phi$  is of fiber type, then as in the case  $[D] = 0$  we get  $X \simeq Z \times \mathbb{P}^1$ . Assume that  $\phi$  is of divisorial type. By inductive hypothesis,  $\phi$  contracts a component of  $[D]$ .

Finally, consider the case when  $\varphi|_S: S \rightarrow Z$  is generically finite of degree 2. Obviously,  $D^{\text{hor}} = S$ . If  $\rho(Z) = 1$ , then all components of  $D^{\text{ver}}$  are numerically proportional and  $\sum_{\text{ver}} d_i \geq 4$ . If  $\dim(W) = 2$ , then  $D^{\text{ver}} \cdot \phi^{-1}(w) \leq 2$  for general

$w \in W$ . Hence  $\sum_{\text{ver}} d_i \geq 4$ , a contradiction. Then by lemmas 3.3 and 3.8,  $\phi$  is divisorial and  $\phi$  must contract  $S$ . We derive a contradiction with Theorem 2.3 for  $(W, \phi(D^{\text{ver}}))$  near  $\phi(S)$ .

Therefore  $\rho(Z) = 2$  and there is a  $K_Z$ -negative extremal contraction  $\psi: Z \rightarrow V$  onto a curve (see Corollary 4.5.3). Let  $\pi: X \xrightarrow{\varphi} Z \xrightarrow{\psi} V$  be the composition map. Clearly, all fibers of  $\pi$  are irreducible. Write  $D = \sum' d_i D_i + \sum'' d_i D_i = D' + D''$ , where  $\sum'$  (respectively  $\sum''$ ) runs through all components  $D_i$  such that  $\pi(D_i) = \text{pt}$  (respectively  $\phi(D_i) = V$ ). Thus,  $S$  is a component of  $D''$  and components of  $D'$  are numerical proportional. Let  $F$  be a general fiber. Then  $\rho(F) = 2$ . Consider the contraction  $\varphi|_F: F \rightarrow \varphi(F)$  and denote  $D''|_F = D|_F$  by  $\Phi = \sum \alpha_i \Phi_i$ . Then  $(F, \Phi)$  is plt and  $K_F + \Phi \equiv 0$ . Clearly, the curve  $S|_F = \lfloor \Phi \rfloor$  is a 2-section and components of  $\Phi - \lfloor \Phi \rfloor$  are fibers of  $\varphi|_F$ . As in the proof of Lemma 3.8 using the fact that  $S|_F$  intersects components of  $\Phi - \lfloor \Phi \rfloor$  twice, one can check  $\sum \alpha_i < 3$ . This yields  $\sum'' d_i < 3$  and  $\sum' d_i > 2$ . Let  $R$  be a  $(K_X + D'')$ -negative extremal ray. Since  $\sum' d_i > 2$  and  $\rho(X) > 2$ ,  $R$  cannot be fiber type. According to Lemma 3.3  $R$  also cannot be flipping type. Therefore  $R$  is divisorial and contract  $S$  to a point. Since  $S$  intersects all components of  $D^{\text{ver}}$ , this contradicts Theorem local. The proof of Proposition 2.4 is finished.

*Concluding remark.* In contrast with the purely log terminal case we have no complete results in the log canonical case. The reason is that the steps of MMP are not so simple. In particular, we can have divisorial contractions which contract components of  $\lfloor D \rfloor$ .

## REFERENCES

- [1] Alexeev V. *Two two-dimensional terminations*, Duke Math. J. **69** (1992) 527–545
- [2] Kawamata Y., Matsuda K. & Matsuki K. *Introduction to the minimal model program*, Adv. Stud. Pure Math. **10** (1987) 283–360
- [3] Kollár J. *Singularities of pairs*, Proc. Symp. Pure Math. **62** (1995) 221–287
- [4] Kollár J. et al. *Flips and abundance for algebraic threefolds*, A summer seminar at the University of Utah, Salt Lake City, 1991. Astérisque. **211** (1992)
- [5] Miyaoka Y. & Mori S. A numerical criterion for uniruledness, Ann. Math. **124** (1986) 65–69
- [6] Mori S. *Threefolds whose canonical bundles are not numerically effective*, Ann. Math. **115** (1982) 133–176
- [7] Iskovskikh V. A. *On the rationality problem for conic bundles*, Math. USSR-Sb. **72** (1992) 105–111
- [8] Iskovskikh V. A. & Prokhorov Yu. G. *Fano Varieties*, Encyclopaedia Math. Sci. **47**, Springer-Verlag (1999)
- [9] Prokhorov Yu. G. *Lectures on complements on log surfaces*, e-print math.AG/9912111
- [10] Sarkisov V. G. *On conic bundle structures*, Math. USSR, Izv. **20** (1983) 355–390
- [11] Shokurov V. V. *3-fold log flips*, Russian Acad. Sci. Izv. Math. **40** (1993) 93–202
- [12] Shokurov V. V. *Complements on surfaces*, e-preprint alg-geom/9711024, to appear in J. Math. Sci.

DEPARTMENT OF MATHEMATICS (ALGEBRA SECTION), MOSCOW LOMONOSOV UNIVERSITY, 117234 MOSCOW, RUSSIA & DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, OH-OKAYAMA, MEGURO, 152 TOKYO, JAPAN  
*E-mail address:* prokhorov@mech.math.msu.su